Turn Over

QP CODE: 24001052

B.Sc DEGREE (CBCS) REGULAR / REAPPEARANCE EXAMINATIONS, MARCH 2024

Sixth Semester

CORE COURSE - MM6CRT01 - REAL ANALYSIS

Common for B.Sc Mathematics Model I, B.Sc Mathematics Model II Computer Science & B.Sc Computer Applications Model III Triple Main

2017 Admission Onwards

25168F10

Time: 3 Hours

Part A

Answer any **ten** questions. Each question carries **2** marks.

- 1. Let f be defind for all $x \in R, x \neq 2$ by $f(x) = \frac{x^2+x-6}{x-2}$. Define f at x = 2 in such a way that f is continuous at that point.
- 2. Give an example of a function $f:[0,1] \rightarrow R$ that is discontinuous at every point of [0,1] but such that |f| is continuous on [0,1].
- 3. Define absolute maximum point and absolute minimum point for f:A
 ightarrow R.
- 4. Is every continuous function differentiable? Justify with proper reasoning or counter example.
- 5. Given that the function $f: R \to R$ defined by $f(x) = x^3 + 2x + 1$ is invertible and let g be its inverse. Find the value of g'(1).
- 6. Define decreasing function with a proper example.
- 7. Define norm of the partition of an interval.
- 8. Test the function of $f(x) = x^{2020} + 2021x$ on [2022, 2023] is Riemann integrable or not.
- 9. Under what circumstances differentiation and Riemann integration are inverse to each other.
- 10. Evaluate $lim(\frac{sinnx}{1+nx})$ for $x \epsilon R, x \ge 0$.
- 11. Show that the sequence of functions f_n defined on R as $f_n(x) = \frac{sin(nx+n)}{n}$ converges uniformly in R.

Page 1/3







Max. Marks: 80



12. Do the limit of a convergent sequence of differentiable functions on an interval [a, b] is differentiable, if not what condition will make the limit function differentiable?

 $(10 \times 2 = 20)$

Part B

Answer any **six** questions. Each question carries **5** marks.

- 13. Define Thomae's function on $(0, \infty)$ and show that it is continuous precisely at the irrational points in $(0, \infty)$.
- 14. State and prove Preservation of Intervals Theorem.
- 15. Let $I \subseteq R$ be an interval and let $f : I \to R$ be monotone on I. Then prove that the set of points $D \subseteq I$ at which f is discontinuous is a countable set .
- ^{16.} Let $f: R \to R$ defined by $f(x) = \begin{cases} x^2, & x \text{ is rational} \\ x, & x \text{ is irrational} \end{cases}$. Prove that f is differentiable at x = 0.
- 17. Derive the inequality $\ x^lpha \leq lpha x + (1-lpha), orall x \geq 0, 0 < lpha < 1$
- 18. Evaluate the limit $\lim_{x o\infty}x^{rac{1}{x}},x\in(0,\infty)$
- 19. Evaluate $\int_{1}^{4} \frac{\sin\sqrt{t}}{\sqrt{t}} dt$.
- 20. Evaluate $\int_{0}^{2} t^{2} (1+t^{3})^{\frac{-1}{2}} dt$.
- 21. Suppose that (f_n) is a sequence of continuous functions on an interval I that converges uniformly on I to a function f. If $(x_n) \subseteq I$ converges to $x_0 \in I$, show that $lim(f_n(x_n)) = f(x_0)$.

(6×5=30)

Part C

Answer any **two** questions.

Each question carries **15** marks.

- 22. (a) Show that a function f is uniformly continuous on the interval (a,b) if and only if it can be defined at the endpoints a and b such that the extended function is continuous on [a,b].(b) State and prove the Continuous Inverse Theorem.
- 23. (a) State and Prove L'Hospital's Rule I
 - (b) Using this, find the following





(i)
$$\lim_{x
ightarrow 0+}rac{ an x-x}{x^3}, x\in(0,rac{\pi}{2})$$

(ii) $\lim_{x
ightarrow 0+}rac{\log\cos x}{x}$

24. (a) Let $f \in \mathcal{R}[a, b]$ and if (\mathcal{P}_n) is any sequence of tagged partitions of [a, b] such that $||\mathcal{P}_n|| \to 0$, prove that $\int_a^b f = lim_n S(f; \mathcal{P})$.

(b) Suppose that f is bounded on [a, b] and that there exists two sequences of tagged partitions (\mathcal{P}_n) and (\mathcal{Q}_n) of [a, b]such that $||\mathcal{P}_n|| \to 0$ and $||\mathcal{Q}_n|| \to 0$, but such that $lim_n S(f; \mathcal{P}_n) \neq lim_n S(f; \mathcal{Q}_n)$. Show that $f \notin \mathcal{R}[a, b]$.

- 25. (a) State and prove the Cauchy Criterion for Riemann integrability of a function $f:[a,b] o\mathbb{R}.$
 - (b) Check the Riemann integrability of Dirichlet function.

(2×15=30)